

# Trichrome Stain, Gomori One-Step, Fast Green - Technical Memo

<b>SOLUTION:</b>	<b>250 ml</b>	<b>500 ml</b>
Trichrome Stain, Gomori One-Step, Fast Green	Part 14021A	Part 14021B

Additionally Needed:

Trichrome, Liver Control Slides	Part 4690	or	Trichrome, Multi-Tissue Control Slides	Part 4693
Xylene, ACS	Part 1445			
Alcohol, Ethyl Denatured, 100%	Part 10841			
Alcohol, Ethyl Denatured, 95%	Part 10842			
Bouin Fluid	Part 1020			
Hematoxylin Stain Set, Weigert Iron	Part 1409			
Acetic Acid 0.5%, Aqueous	Part 100121			
Coplin Jar, Plastic	Part 5184 (for microwave modification)			

**For storage requirements and expiration date refer to individual product labels.**

**APPLICATION:**

Newcomer Supply Trichrome Stain, Gomori One-Step, Fast Green procedure, with included microwave modification, uses a one-step solution combining a plasma stain and a connective tissue stain to differentially demonstrate collagen and muscle fibers.

**METHOD:**

**Fixation:** Formalin 10%, Phosphate Buffered (Part 1090)

**Technique:** Paraffin sections cut at 4 microns

a. See Procedure Note #1.

**Solutions:** All solutions manufactured by Newcomer Supply, Inc.

All Newcomer Supply stain procedures are designed to be used with Coplin jars filled to 40 ml following the staining procedure.

**PRESTAINING PREPARATION:**

1. If necessary, heat dry tissue sections/slides in oven.
2. Preheat Bouin Fluid (1020) to 56-60°C in oven or water bath. **(Skip if using overnight method or microwave procedure.)**

**STAINING PROCEDURE:**

3. Deparaffinize sections thoroughly in three changes of xylene, 3 minutes each. Hydrate through two changes each of 100% and 95% ethyl alcohols, 10 dips each. Wash well with distilled water.
  - a. See Procedure Notes #2 and #3.
4. Mordant in preheated Bouin Fluid (Step #2) for 1 hour at 56-60°C or overnight at room temperature. Cool at room temperature for 5-10 minutes.
  - a. Skip Step #4 if tissue was originally Bouin fixed.

**Microwave Modification:** See Procedure Note #4.

b. Place slides in a plastic Coplin jar containing Bouin Fluid and microwave for 5 minutes at 60°C. Allow slides to sit an additional 10 minutes in solution.

5. Wash well in running tap water; rinse in distilled water.
6. Prepare fresh Weigert Iron Hematoxylin (1409); combine and mix well.
  - a. Solution A: Ferric Chloride, Acidified 20 ml
  - b. Solution B: Hematoxylin 1%, Alcoholic 20 ml
7. Stain slides in fresh Weigert Iron Hematoxylin for 10 minutes.
8. Wash in running tap water for 10 minutes; rinse in distilled water.
  - a. See Procedure Note #5.
9. Stain in Trichrome Stain, Gomori One-Step, Fast Green for 20 minutes.
10. Differentiate in Acetic Acid 0.5%, Aqueous (100121) for 2 minutes.

11. Rinse quickly in distilled water.
12. Dehydrate in two changes each of 95% and 100% ethyl alcohol. Clear in three changes of xylene, 10 dips each; coverslip with compatible mounting medium.

**RESULTS:**

Collagen and mucin	Green
Muscle fibers, cytoplasm and keratin	Red
Nuclei	Blue/black

**PROCEDURE NOTES:**

1. Using ammonium hydroxide to soak/face tissue blocks will alter the pH of tissue sections and diminish trichrome staining.
2. Drain slides after each step to prevent solution carry over.
3. Do not allow sections to dry out at any point during procedure.
4. The suggested microwave procedure has been tested at Newcomer Supply. This procedure is a guideline and techniques should be developed for your laboratory.
5. If Weigert Iron Hematoxylin is not completely washed from tissue sections, nuclear and cytoplasmic staining may be compromised.
6. If using a xylene substitute, closely follow the manufacturer's recommendations for deparaffinization and clearing steps.

**REFERENCES:**

1. Brown, Richard. *Histologic Preparations: Common Problems and Their Solutions*. Northfield, Ill.: College of American Pathologists, 2009. 95-101.
2. Carson, Freida L., and Christa Hladik. *Histotechnology: A Self-Instructional Text*. 3rd ed. Chicago, Ill.: American Society of Clinical Pathologists, 2009. 165-166.
3. Sheehan, Dezna C., and Barbara B. Hrapchak. *Theory and Practice of Histotechnology*. 2nd ed. St. Louis: Mosby, 1980. 191-192.
4. Vacca, Linda L. *Laboratory Manual of Histochemistry*. New York: Raven Press, 1985. 308-310.
5. Modifications developed by Newcomer Supply Laboratory.